

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications, Department of Psychology

Psychology, Department of

2015

Biomechanics of head impacts associated with diagnosed concussion in female collegiate ice hockey players

Bethany J. Wilcox

The Warren Alpert Medical School of Brown University and Rhode Island Hospital

Jonathan G. Beckwith

Simbex, Lebanon, NH

Richard M. Greenwald

Dartmouth College, richard.m.greenwald@dartmouth.edu

Neha P. Raukar

The Warren Alpert Medical School of Brown University

Jeffrey J. Chu

Simbex, Lebanon, NH

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/psychfacpub>



Part of the [Psychology Commons](#)

Wilcox, Bethany J.; Beckwith, Jonathan G.; Greenwald, Richard M.; Raukar, Neha P.; Chu, Jeffrey J.; McAllister, Thomas W.; Flashman, Laura A.; Maerlender, Arthur C.; Duhaime, Ann-Christine; and Crisco, Joseph J., "Biomechanics of head impacts associated with diagnosed concussion in female collegiate ice hockey players" (2015). *Faculty Publications, Department of Psychology*. 713.

<http://digitalcommons.unl.edu/psychfacpub/713>

This Article is brought to you for free and open access by the Psychology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Psychology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Bethany J. Wilcox; Jonathan G. Beckwith; Richard M. Greenwald; Neha P. Raukar; Jeffrey J. Chu; Thomas W. McAllister; Laura A. Flashman; Arthur C. Maerlender; Ann-Christine Duhaime;; and Joseph J. Crisco

Biomechanics of head impacts associated with diagnosed concussion in female collegiate ice hockey players

Bethany J. Wilcox,¹ Jonathan G. Beckwith,² Richard M. Greenwald,^{2,3} Neha P. Raukar,⁴
Jeffrey J. Chu,² Thomas W. McAllister,⁵ Laura A. Flashman,⁶ Arthur C. Maerlender,⁷
Ann-Christine Duhaime,⁸ Joseph J. Crisco¹

¹ Bioengineering Laboratory, Department of Orthopaedics,
The Warren Alpert Medical School of Brown University and Rhode Island Hospital, Providence, RI
² Simbex, Lebanon, NH

³ Thayer School of Engineering, Dartmouth College, Hanover, NH

⁴ Department of Emergency Medicine, The Warren Alpert Medical School of Brown University, Providence, RI

⁵ Department of Psychiatry, Indiana University School of Medicine, Indianapolis, IN

⁶ Department of Psychiatry, Geisel School of Medicine at Dartmouth, Lebanon, NH

⁷ Center for Brain Biology and Behavior, University of Nebraska–Lincoln, Lincoln, NE

⁸ Pediatric Neurosurgery, Massachusetts General Hospital, Boston, MA

Corresponding author — B.J. Wilcox, Bioengineering Laboratory, Department of Orthopaedics,
Alpert Medical School of Brown University and Rhode Island Hospital, CORO West, Suite 404, 1 Hoppin Street,
Providence, RI 02903, USA; tel 401 263-7080, fax 401 444-4418, email Bethany_Wilcox@brown.edu

Abstract

Epidemiological evidence suggests that female athletes may be at a greater risk of concussion than their male counterparts. The purpose of this study was to examine the biomechanics of head impacts associated with diagnosed concussions in a cohort of female collegiate ice hockey players. Instrumented helmets were worn by 58 female ice hockey players from 2 NCAA programs over a three year period. Kinematic measures of single impacts associated with diagnosed concussion and head impact exposure on days with and without diagnosed concussion were evaluated. Nine concussions were diagnosed. Head impact exposure was greater in frequency and magnitude on days of diagnosed concussions than on days without diagnosed concussion for individual athletes. Peak linear accelerations of head impacts associated with diagnosed concussion in this study are substantially lower than those previously reported in male athletes, while peak rotational accelerations are comparable. Further research is warranted to determine the extent to which female athletes' biomechanical tolerance to concussion injuries differs from males.

Keywords: Impact biomechanics, Concussion, Female, Hockey

1. Introduction

Epidemiological evidence suggests that female athletes may be at a greater risk of concussion than their male counterparts (Dick, 2009). Research conducted in sports where males and females participate at the same level, such as ice hockey, has shown that female athletes sustain concussions at a higher rate (Agel and Harvey, 2010 and Dick, 2009). It has been reported that females experience a greater number of symptoms associated with concussions and that these symptoms are often more severe (Broshek et al., 2005 and Colvin et al., 2009). Recent studies have also suggested that females may be at a greater risk of post-concussive syndrome, where the acute symptoms associated with concussion injuries become chronic (Preiss-Farzanegan et al., 2009). Several sex-specific characteristics have been proposed as the rationale for these differences, including physiological differences and

psychological factors, but the exact reasons remain unclear. While it has been generally accepted that the mechanism of concussion injury is related to head acceleration (McCroly et al., 2009), the exact relationship between the biomechanics of head impacts, sex, and clinical outcome of concussion is unknown.

Recent studies have focused on understanding the biomechanics of head impacts sustained in contact sports by utilizing an accelerometer based head impact monitoring device, the Head Impact Telemetry (HIT) System (Simbex, Lebanon, NH). The HIT System allows researchers to monitor and record head impacts sustained by individual helmeted athletes during play. Utilizing this system, researchers have quantified head impact exposure, a multifactorial term including frequency, magnitude and location on the helmet of head impacts, for individual athletes at different levels of play in football and ice hockey (Broglia et al., 2009, Crisco et al., 2011, Mihalik et al., 2008 and Wilcox et al., 2013).

Data from the HIT System has also been used to evaluate the biomechanics of head impacts associated with diagnosed concussions in high school and collegiate football players (Beckwith et al., 2013, Funk et al., 2012, Guskiewicz et al., 2007 and Rowson et al., 2012). Beckwith et al. (2013) reported that individual players sustained more head impacts and impacts of greater severity on days of diagnosed concussion than on days without. While these studies have provided valuable insights into the biomechanics of impacts associated with diagnosed concussion in male football players, quantification and analysis of head impacts associated with diagnosed concussion in female athletes are currently lacking.

The purpose of this study was to evaluate the biomechanics of head impacts associated with diagnosed concussions in a study population of female collegiate ice hockey players.

2. Methods

Over a three year period, 58 female ice hockey players from 2 National Collegiate Athletic Association (NCAA) athletic programs (Brown University and Dartmouth College) participated in this observational study after informed consent was obtained with institutional review board approval. Players participated during multiple years; 19, 20, and 19 players participated in one, two, and three seasons respectively. This participant turnover is typical of fluctuations on collegiate athletic teams.

All participants wore instrumented helmets (Easton S9, Van Nuys, CA; Head Impact Telemetry (HIT) System, Simbex, Lebanon, NH) to collect the magnitude of head impacts sustained during practices and games. These helmets measured and recorded biomechanical data from head impacts including linear and rotational acceleration of the center of gravity of the head. The instrumented helmets each consist of six single-axis accelerometers arranged tangentially to the head and mounted elastically within the liners of helmets to maintain contact with the head, thus isolating the accelerometers from shell vibrations (Brainard et al., 2012, Manoogian et al., 2006 and Mihalik et al., 2008). System design, processing methods, accuracy, and data reduction techniques have been previously described in detail (Chu et al., 2006, Crisco et al., 2004, Duma et al., 2005, Greenwald et al., 2008 and Mihalik et al., 2008). Acceleration data is collected at 1 kHz, time stamped and stored (up to 100 impacts in static memory). Data is then transmitted by radiofrequency telemetry to a computer and entered into a secure database.

For this study, concussions were defined as an alteration in mental status, as reported or observed by the player or medical staff, resulting from a blow to the head. Diagnosis of concussion was made by the team physician or certified athletic trainer (ATC) at each respective institution. For each diagnosed concussion, suspected date, time, and symptom presentation within 72 h of the injury were documented. Anecdotal descriptions of the events surrounding injury were also collected to further aid in corroborating data from the instrumented helmets with diagnosed concussions.

To evaluate the biomechanics of impacts associated with diagnosed concussion, two different approaches were taken: (1) the examination of single impacts linked to injury, and (2) the cumulative analysis of all impacts sustained on days of injury. Single impacts were identified when a concussion event was attributed by the player or observers to a specific identifiable impact. Impact magnitude variables included peak linear acceleration (g), peak rotational acceleration (rad/s^2), and HITsp (Crisco et al., 2004 and Greenwald et al., 2008). HITsp is a weighted measure of head impact severity that includes linear and rotational acceleration, impact duration, and impact location (Greenwald et al., 2008). For the cumulative analysis, head impact exposure was computed using a single measure of impact frequency, the total number of impacts per day, for individual players on days with and without diagnosed concussion (Beckwith et al., 2013). Each individual player's distribution of peak linear acceleration (g), peak rotational acceleration (rad/s^2), and HITsp per day was quantified by the 50th and 95th percentile value of all impacts per day. Results are expressed as median values and 25–75% interquartile range of those percentile values.

2.1 Statistical analysis

Head impact frequency and magnitude variables were found to be non-normally distributed (Shapiro–Wilk test, $p < 0.05$). Therefore, to evaluate differences in these measures for days with and without concussion, Wilcoxon sign-ranked tests for pairs were used. Statistical analyses were performed using SigmaPlot 12.0 (Systat Software, Chicago, IL).

3. Results

There were a total of nine diagnosed concussions and approximately half of these injuries (4/9) were associated with an identifiable single impact. The average (\pm standard deviation) peak linear acceleration, peak rotational acceleration, and HITsp of these impacts were $43.0 \pm 11.5g$, $4030 \pm 1435 \text{ rad/s}^2$, and 25.6 ± 4.8 , respectively (Table 1).

Significant differences were observed in the frequency and magnitude of head impacts that players sustained on days with and without diagnosed concussion. Players sustained a greater number of head impacts ($p = 0.023$) on days of diagnosed concussion than on days without (Fig. 1). The 50th and 95th percentile values of peak linear acceleration and peak rotational acceleration were significantly higher on days with diagnosed concussion than on days without (Table 2). Ninety-fifth percentile HITsp was also significantly higher on days with diagnosed concussion than on days without.

4. Discussion

The purpose of this study was to examine the biomechanics of head impacts associated with diagnosed concussion in a study population of female collegiate ice hockey players. Data was collected over a three year span from approximately 60 athletes. While 15% of these athletes were diagnosed with a concussion during the course of the study, this yielded only nine concussion cases for analysis. Herein, we report and discuss our findings with full disclosure of the limitations of such a small dataset.

Substantial research has focused on single impacts associated with concussion injury with the objective of identifying a biomechanical threshold for the injury. A study in a large cohort of high school and collegiate football players reported an average peak linear and peak rotational acceleration of $112.1 \pm 35.4g$ and $4253 \pm 2287 \text{ rad/s}^2$ for single impacts associated with diagnosed concussion (Beckwith et al., 2013). Other studies in football have reported peak linear accelerations associated with concussions ranging from 60.51 to 168.71g (Broglia et al., 2012 and Guskiewicz et al., 2007). In a study of collegiate football players, the average linear acceleration of concussive impacts was $145 \pm 35g$, and a risk analysis resulted in a threshold of 100 g as a cutoff for identifying potential concussions (Funk et al., 2012). Comparable to a study in a larger cohort of athletes, approximately 50% (4/9) of the diagnosed concussions in our study were associated with a single identifiable impact (Duhaime et al., 2012). The average peak linear acceleration of the impacts associated with diagnosed concussions in the current study ($43 \pm 11.5g$) is substantially lower than any values reported for football players in the literature. If the recommended cutoff for collegiate football players (100 g) was used to monitor this group of female athletes, not one of the impacts that were associated with diagnosed concussions would have reached this threshold. These findings suggest that biomechanical thresholds may need to be sex-specific. Interestingly, the average peak rotational acceleration of these impacts ($4029.5 \pm 1243 \text{ rad/s}^2$) is comparable to those previously reported in male football players.

Table 1. Peak linear acceleration, peak rotational acceleration, and HITsp for single impacts ($n=4$) associated with diagnosed concussions in a cohort of female collegiate ice hockey players.

Peak linear acceleration (g)	Peak rotational acceleration (rad/s^2)	HITsp
53.3	1959	18.8
30.4	4945	27.0
36.1	5045	26.5
52.2	4169	30.1

While studies have shown that both linear and rotational accelerations of the head play a role in concussion, the relative contributions of these kinematic components to this complex injury remain unknown (Guskiewicz and Mihalik, 2011). In this study, peak linear accelerations associated with diagnosed concussion, which has been shown to be the most predictive impact severity metric in football (Beckwith et al., 2013), suggest that females have a lower biomechanical tolerance to concussion injuries than their male counterparts. However, when considering peak rotational acceleration alone, the measures recorded from female and male athletes are remarkably similar which could support a similar tolerance to rotational acceleration. While it is well established that linear and rotational head accelerations are correlated, particularly in helmeted contact sports (Pellman et al., 2003, Rowson et al., 2012 and Rowson et al., 2009), the exact relationship for individual impacts depends upon several factors including impact location, anticipation of impact and, accordingly, neck strength. Given these factors, it is possible that this relationship differs between males and females.

Similar to previous studies, we found that head impact exposure on days of diagnosed concussion is higher for individual athletes than on days without diagnosed concussion, in both frequency and magnitude (Beckwith et al., 2013). This supports previous assertions that diagnosis of concussion is related to head impact exposure (Beckwith et al., 2013 and Crisco and Greenwald, 2011). Several measures of head impact exposure on days of diagnosed concussions for the female collegiate hockey players in the current study, however, were substantially lower than the same variables previously reported for male football players (Beckwith et al., 2013). This further supports the postulation that female athletes may have a differing biomechanical tolerance to concussions than their male counterparts.

Several sex-specific characteristics have been proposed to explain differences between males and females in concussion injuries including physiological differences, such as anthropometrics and hormone regulation (Bramlett and Dietrich, 2001 and Stein and Hoffman, 2003), and psychological factors, including speculation that females tend to be more honest than males in reporting injuries (Covassin et al., 2012 and Dick, 2009). Whether this reflects actual increased tissue damage or greater symptom intensity for a given biomechanical input, increased concern or focus on potential consequences of injury, or some other factor remains speculative.

This study has several limitations. While approximately 15% of the athletes in the study were diagnosed with a concussion, this accounts for a relatively small sample size of concussion cases ($n = 9$). Additionally, a limitation in evaluating single impacts associated with diagnosed concussions is the inherent error associated with the use of any measurement device, in this case the HIT System. In appreciation of these limitations, the values reported in this short communication should be interpreted carefully. While several biomechanical measures of head impacts associated with diagnosed concussion in female athletes were substantially lower than data previously reported in male athletes collected with




Fig. 1. Median [25–75%] of the 95th percentile peak linear acceleration as a function of the median [25–75%] frequency of impacts per day. Filled markers represent days with diagnosed concussion; unfilled markers represent days without diagnosed concussion.

similarly instrumented helmets with presumably similar limitations, further investigation is necessary to draw any conclusions. To address the concerns associated with evaluating a limited dataset of single impacts, we analyzed head impact exposure cumulatively for the days that players were and were not diagnosed with concussion injuries. However, an additional limitation exists in that it is not known whether a cumulative measure over a longer period of time would be more appropriate than limiting the analysis to a single day.

Another important consideration in this study is that diagnosis of concussion was represented as a binary state and did not acknowledge that there exists a spectrum of concussive injuries (Duhaime et al., 2012). The diagnosed concussions presented in this study exhibited a spectrum of signs and symptoms that may represent very different injuries. Moreover, it is possible that the diagnosed concussions in the female ice hockey players were of a lower severity than those previously reported in male athletes.

We reported the biomechanics of head impacts associated with nine diagnosed concussions in a study population of female ice hockey players. Head impact exposure was greater in frequency and magnitude on days of diagnosed concussions than on days without diagnosed concussion for individual athletes. Peak linear accelerations of impacts associated with diagnosed concussion were found to be substantially lower than those previously reported in male athletes, while rotational accelerations were comparable. Further research is warranted to determine the extent to which female athletes' biomechanical tolerance to concussion injuries differs from males.

Table 2. Magnitudes, reported as median (25–75% interquartile range), of head impacts sustained on days with diagnosed concussion were more severe than on days without diagnosed concussion.

	Percentile	Days w/diagnosed concussion	Days w/out diagnosed concussion	p-value
Peak linear acceleration(g)	50th	21.9 (17.4–22.8)	15.3 (15.1–15.6)	*p=0.001
	95th	36.1 (30.4–51.6)	20.7 (18.6–22.0)	*p=0.001
Peak rotational acceleration (rad/s ²)	50th	2046 (1676–2202)	1249 (1169–1397)	*p=0.047
	95th	3491 (2411–4169)	1769 (1621–2123)	*p=0.015
HITsp	50th	14.7 (13.5–18.4)	13.5 (13.4–13.7)	p=0.241
	95th	23.2 (17.9–26.5)	15.4 (14.8–16.2)	*p=0.012

Conflict of interest disclosure — Joseph J. Crisco, Richard M. Greenwald, and Simbex have a financial interest in the instruments (HIT System, Sideline Response System (Riddell, Inc.)) that were used to collect the biomechanical data reported in this study.

Acknowledgments — Research reported in this publication was supported by the National Institute of Child Health and Human Development, the National Institute of Neurological Disorders and Stroke, and the National Institute of General Medical Sciences at the National Institute of Health under award numbers R01HD048638, R01NS055020, R25GM083270 and R25GM083270-S1 and the National Operating Committee on Standards for Athletic Equipment (NOCSAE). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or NOCSAE. We gratefully acknowledge and thank the engineering team at Simbex for all of their technical support. We would like to thank Lindley Brainard and Wendy Chamberlin at Simbex for their role in data collection and clinical coordination. We would also like to thank Russell Fiore, M.Ed., A.T.C., Emily Burmeister M.S., A.T.C., and Brian Daigneault, MS, A.T.C. at Brown University; Jeff Frechette, A.T.C., and Tracey Poro, A.T.C. at Dartmouth College Sports Medicine and Mary Hynes, R.N., M.P.H. at Dartmouth Medical School for their support on this project.

References

- Agel and Harvey, 2010 □ J. Agel, E.J. Harvey, A 7-year review of men's and women's ice hockey injuries in the NCAA, *Can. J. Surg.*, 53 (2010), pp. 319–323.
- Beckwith et al., 2013 □ J.G. Beckwith, R.M. Greenwald, J.J. Chu, J.J. Crisco, S. Rowson, S.M. Duma, S.P. Broglio, T.W. McAllister, K.M. Guskiewicz, J.P. Mihalik, S. Anderson, B. Schnebel, P.G. Brolinson, M.W. Collins, Head impact exposure sustained by football players on days of diagnosed concussion, *Med. Sci. Sports Exerc.*, 45 (2013), pp. 737–746.
- Brainard et al., 2012 □ L.L. Brainard, J.G. Beckwith, J.J. Chu, J.J. Crisco, T.W. McAllister, A.-C. Duhaime, A.C. Maerlender, R.M. Greenwald, Gender differences in head impacts sustained by collegiate ice hockey players, *Med. Sci. Sports Exerc.*, 44 (2012), pp. 297–304.
- Bramlett and Dietrich, 2001 □ H.M. Bramlett, W.D. Dietrich, Neuropathological protection after traumatic brain injury in intact female rats versus males or ovariectomized females, *J. Neurotrauma*, 18 (2001), pp. 891–900.
- Broglio et al., 2012 □ S.P. Broglio, J.T. Eckner, J.S. Kutcher, Field-based measures of head impacts in high school football athletes, *Curr. Opin. Pediatr.*, 24(6) (2012), pp. 702–708.
- Broglio et al., 2009 □ S.P. Broglio, J.J. Sosnoff, S. Shin, X. He, C. Alcaraz, J. Zimmerman, Head impacts during high school football: a biomechanical assessment, *J. Athl. Train.*, 44 (2009), pp. 342–349.
- Broshek et al., 2005 □ D.K. Broshek, T. Kaushik, J.R. Freeman, D. Erlanger, F. Webbe, J.T. Barth, Sex differences in outcome following sports-related concussion, *J. Neurosurg.*, 102 (2005), pp. 856–863.
- Chu et al., 2006 □ J.J. Chu, J.G. Beckwith, J.J. Crisco, R.M. Greenwald, A novel algorithm to measure linear and rotational head acceleration using single-axis accelerometers, *J. Biomech.*, 39 (2006), p. S534.
- Colvin et al., 2009 □ A.C. Colvin, J. Mullen, M.R. Lovell, R.V. West, M.W. Collins, M. Groh, The role of concussion history and gender in recovery from soccer-related concussion, *Am. J. Sports Med.*, 37 (2009), pp. 1699–1704.
- Covassin et al., 2012 □ T. Covassin, R.J. Elbin, W. Harris, T. Parker, A. Kontos, The role of age and sex in symptoms, neurocognitive performance, and postural stability in athletes after concussion, *Am. J. Sports Med.*, 40 (2012), pp. 1303–1312.
- Crisco et al., 2004 □ J.J. Crisco, J.J. Chu, R.M. Greenwald, An algorithm for estimating acceleration magnitude and impact location using multiple nonorthogonal single-axis accelerometers, *J. Biomech. Eng.*, 126 (2004), pp. 849–854.
- Crisco and Greenwald, 2011 □ J.J. Crisco, R.M. Greenwald, Let's get the head further out of the game: a proposal for reducing brain injuries in helmeted contact sports, *Curr. Sports Med. Rep.*, 10 (2011), pp. 7–9.
- Crisco et al., 2011 □ J.J. Crisco, B.J. Wilcox, J.G. Beckwith, J.J. Chu, A.-C. Duhaime, S. Rowson, S.M. Duma, A.C. Maerlender, T.W. McAllister, R.M. Greenwald, Head impact exposure in collegiate football players, *J. Biomech.*, 44 (2011), pp. 2673–2678.
- Dick, 2009 □ R.W. Dick, Is there a gender difference in concussion incidence and outcomes?, *Br. J. Sports Med.*, 43 (Suppl 1) (2009), pp. Si46–Si50.
- Duhaime et al., 2012 □ A.-C. Duhaime, J.G. Beckwith, A.C. Maerlender, T.W. McAllister, J.J. Crisco, S.M. Duma, P.G. Brolinson, S. Rowson, L.A. Flashman, J.J. Chu, R.M. Greenwald, Spectrum of acute clinical characteristics of diagnosed concussions in college athletes wearing instrumented helmets: clinical article, *J. Neurosurg.*, 117 (2012), pp. 1092–1099.
- Duma et al., 2005 □ S.M. Duma, S.J. Manoogian, W.R. Bussone, P.G. Brolinson, M.W. Goforth, J.J. Donnenwerth, R.M. Greenwald, J.J. Chu, J.J. Crisco, Analysis of real-time head accelerations in collegiate football players, *Clin. J. Sport Med.*, 15 (2005), pp. 3–8.
- Funk et al., 2012 □ J.R. Funk, S. Rowson, R.W. Daniel, S.M. Duma, Validation of concussion risk curves for collegiate football players derived from HITS data, *Ann. Biomed. Eng.*, 40 (2012), pp. 79–89.
- Greenwald et al., 2008 □ R.M. Greenwald, J.T. Gwin, J.J. Chu, J.J. Crisco, Head impact severity measures for evaluating mild traumatic brain injury risk exposure, *Neurosurgery*, 62 (2008), pp. 789–798.
- Guskiewicz and Mihalik, 2011 □ K.M. Guskiewicz, J.P. Mihalik, Biomechanics of sport concussion: quest for the elusive injury threshold, *Exerc. Sport Sci. Rev.*, 39 (2011), pp. 4–11.
- Guskiewicz et al., 2007 □ K.M. Guskiewicz, J.P. Mihalik, V. Shankar, S.W. Marshall, D.H. Crowell, S.M. Oliaro, M.F. Ciocca, D.N. Hooker, Measurement of head impacts in collegiate football players: relationship between head impact biomechanics and acute clinical outcome after concussion, *Neurosurgery*, 61 (2007), pp. 1244–1252 discussion 1252–1253.
- Manoogian et al., 2006 □ S. Manoogian, D. McNeely, S. Duma, G. Brolinson, R. Greenwald, Head acceleration is less than 10 percent of helmet acceleration in football impacts, *Biomed. Sci. Instrum.*, 42 (2006), pp. 383–388.
- McCrory et al., 2009 □ P. McCrory, W. Meeuwisse, K. Johnston, J. Dvorak, M. Aubry, M. Molloy, R. Cantu, Consensus statement on concussion in sport: the 3rd international conference on concussion in sport held in Zurich, *J. Athl. Train.*, 44 (2009), pp. 434–448.
- Mihalik et al., 2008 □ J.P. Mihalik, K.M. Guskiewicz, J.A. Jeffries, R.M. Greenwald, S.W. Marshall, Characteristics of head impacts sustained by youth ice hockey players, *ImechE*, 222 (2008), pp. 45–52.
- Pellman et al., 2003 □ E.J. Pellman, D.C. Viano, A.M. Tucker, I.R. Casson, J.F. Waeckerle, Concussion in professional football: reconstruction of game impacts and injuries, *Neurosurgery*, 53 (2003), pp. 799–812 discussion 812–814.
- Preiss-Farzanegan et al., 2009 □ S.J. Preiss-Farzanegan, B. Chapman, T.M. Wong, J. Wu, J.J. Bazarian, The relationship between gender and postconcussion symptoms after sport-related mild traumatic brain injury, *PM R*, 1 (2009), pp. 245–253.
- Rowson et al., 2009 □ S. Rowson, G. Brolinson, M. Goforth, D. Dietter, S. Duma, Linear and angular head acceleration measurements in collegiate football, *J. Biomech. Eng.*, 131 (2009), p. 061016.
- Rowson et al., 2012 □ S. Rowson, S.M. Duma, J.G. Beckwith, J.J. Chu, R.M. Greenwald, J.J. Crisco, P.G. Brolinson, A.-C. Duhaime, T.W. McAllister, A.C. Maerlender, Rotational head kinematics in football impacts: an injury risk function for concussion, *Ann. Biomed. Eng.*, 40 (2012), pp. 1–13.
- Stein and Hoffman, 2003 □ D.G. Stein, S.W. Hoffman, Estrogen and progesterone as neuroprotective agents in the treatment of acute brain injuries, *Pediatr. Rehabil.*, 6 (2003), pp. 13–22.
- Wilcox et al., 2013 □ B.J. Wilcox, J.G. Beckwith, R.M. Greenwald, J.J. Chu, T.W. McAllister, L.A. Flashman, A.C. Maerlender, A.C. Duhaime, J.J. Crisco, Head impact exposure in male and female collegiate ice hockey players, *J. Biomech.*, 47(1) (2013), pp. 109–114.